

The Effective Dimension of Cones and Degrees

Jan Reimann

Institut für Informatik, Universität Heidelberg



15.11.2005

Overview

- 1 Effective Hausdorff Dimension
- 2 The Dimension of Cone and Degrees
- 3 A Wtt Lower Cone of Non-Integral Dimension
- 4 The Turing Case

Outer Measures

The Caratheodory Method

- A **premeasure** is a function $\rho : 2^{<\omega} \rightarrow \mathbb{R}_0^+ \cup \{\infty\}$
- One can obtain an **outer measure** μ_ρ from ρ by letting

$$\mu_\rho(\mathcal{X}) = \inf \left\{ \sum_i \rho(x_i) : \bigcup_i [x_i] \supseteq \mathcal{X} \right\}.$$

- $\mu = \mu_\rho$ is a countably subadditive, monotone set function. Restriction to sets \mathcal{A} which satisfy

$$(\forall \mathcal{Y}) \mu(\mathcal{Y}) = \mu(\mathcal{Y} \cap \mathcal{A}) + \mu(\mathcal{Y} \setminus \mathcal{A}),$$

yields the **measurable sets**.

- The measurable sets form a **σ -algebra**, and μ is an additive set function on this σ -algebra.

Effectivizing Measures

- Let ρ be a computable premeasure, with μ_ρ the induced outer measure.

Definition

A set $\mathcal{X} \subseteq 2^\omega$ has **effectively μ_ρ -measure zero** if there exists a uniformly computable sequence (C_n) of sets of strings such that for all n ,

$$\mathcal{X} \subseteq \bigcup_{\sigma \in C_n} [\sigma] \quad \text{and} \quad \sum_{\sigma \in C_n} \rho(\sigma) \leq 2^{-n}.$$

Hausdorff Measures and Hausdorff Dimension

- Hausdorff measures \mathcal{H}^s arise from the premeasures $\rho(\sigma) = 2^{-|\sigma|^s}$, $s \geq 0$.
- It is obvious that $\mathcal{H}^s \mathcal{X} = 0$ implies $\mathcal{H}^t \mathcal{X} = 0$ for all $t > s$.

Definition

The **Hausdorff dimension** of \mathcal{X} is defined as

$$\dim_{\text{H}} \mathcal{X} = \inf\{s \geq 0 : \mathcal{H}^s \mathcal{X} = 0\}.$$

Effective Hausdorff Dimension

Definition

The effective Hausdorff dimension of \mathcal{X} is defined as

$$\dim_{\mathcal{H}}^1 \mathcal{X} = \inf\{s \in \mathbb{Q}_0^+ : \mathcal{X} \text{ is effectively } \mathcal{H}^s\text{-null}\}.$$

[Lutz 2000]

- There are single reals of non-zero dimension: every Martin-Löf random real has dimension one.
- Effective dimension has an important stability property:

$$\dim_{\mathcal{H}}^1 \mathcal{X} = \sup\{\dim_{\mathcal{H}}^1\{A\} : A \in \mathcal{X}\}.$$

[Lutz 2000]

Effective Hausdorff Dimension

Effective dimension as degree of randomness

Theorem

For every real A ,

$$\dim_{\text{H}}^1 A = \liminf_{n \rightarrow \infty} \frac{K(A \upharpoonright_n)}{n} =: \underline{K}(A).$$

[Ryabko 1984; Mayordomo 2002]

Effective Hausdorff Dimension

The three basic examples

- Given $0 < r < 1$ rational, let $Z_r = \{\lfloor n/r \rfloor : n \in \mathbb{N}\}$. Given a Martin-Löf random set X , define X_r by

$$X_r(m) = \begin{cases} X(n) & \text{if } m = \lfloor n/r \rfloor, \\ 0 & \text{otherwise.} \end{cases}$$

Then $\dim_{\mathbb{H}}^1 X_r = r$.

- Let μ_p be a Bernoulli (“coin-toss”) measure with bias $p \in \mathbb{Q} \cap [0, 1]$, and let B be Martin-Löf random with respect to μ_p . Then

$$\dim_{\mathbb{H}}^1 B = H(\mu_p) := -[p \log p + p \log(1 - p)].$$

Effective Hausdorff Dimension

The three basic examples

- Let U be a universal, prefix-free machine. Given a computable real number $0 < s \leq 1$, the binary expansion of the real number

$$\Omega^{(s)} = \sum_{\sigma \in \text{dom}(U)} 2^{-\frac{|\sigma|}{s}}$$

has effective dimension s [Tadaki 2002]. (Note that $\Omega^{(1)}$ is just Chaitin's Ω .)

Effective Hausdorff Dimension

The basic examples imply genuine random content

- Each of the three examples actually computes a Martin-Löf random real.
- This is obvious for the “diluted” sequence.
- For computable Bernoulli measures, one may use [Von-Neumann’s trick](#) to turn a biased random real into a uniformly distributed random real. More generally, Levin (1970) and Kautz (1991) have shown that any real which is random with respect to computable measure computes a Martin-Löf random real.
- $\Omega^{(s)}$ computes a fixed-point free function. It is a left-computable real, and hence it follows from the [Arslanov completeness criterion](#) that $\Omega^{(s)}$ is Turing complete (and thus T-equivalent to a Martin-Löf random real).

The Dimension Problem

Are there “genuine” reals of non-integral dimension?

- The stability property implies that the Turing lower cone of each of the three examples has effective dimension 1.

Question

Are there any Turing lower cones of non-integral dimension?

- This is an **open problem**. Any such lower cone would come from a real of non-integral dimension for which it is not possible to extract some content of higher degree of randomness effectively.

Upper Cones

Upper cones always have maximal dimension

- For upper cones, the situation is quite clear.
- It is known that the Turing upper cone of a real has Lebesgue measure zero unless the real is computable [Sacks 1963].

Theorem

For any real A , the many-one upper cone of A has (classical) Hausdorff dimension 1.

Lower Cones and Degrees

- The dimension of a lower cone and a degree coincide.
- This follows from the **sparse coding technique**: Given two reals $A \leq_r B$, choose a computable real R of density $\lim_n |R \cap \{0, \dots, n-1\}|/n = 1$, and let C equal A on R and B on the complement of R .
- C will be r -equivalent to B and be of the **same dimension as A** . It follows that the dimension of the degree and the lower cone of a set coincide.

Many-One Reducibility

Theorem

Let μ_p be a computable Bernoulli measure with bias p . If A is μ_p -random, then

$$B \leq_m A \Rightarrow \dim_{\text{H}}^1 B \leq H(\mu_p).$$

[Reimann and Terwijn 2004]

- **Proof.** Given an m -reduction f , define $F = \{n : (\forall m < n) f(m) \neq f(n)\}$, so F is the set of all positions of B , where an instance of A is queried for the first time.
- F induces a **Kolmogorov-Loveland place selection rule**. If A is μ_p -random, this selection rule will yield a new sequence with the same limit frequency as A .

Weaker Reducibilities

- This technique does not extend to weaker reducibilities, since for **Bernoulli measures** the Levin-Kautz result holds for a **total Turing reduction**.
- Stephan (2005) was able to construct wtt-lower cone of non-integral effective dimension in a relativized world:
There is a real A and an oracle B such that

$$1/3 \leq \dim_H^B \{D : D \leq_{\text{wtt}}^B A\} \leq 1/2.$$

A Wtt Lower Cone of Non-Integral Dimension

The result

Theorem

For each rational α , $0 \leq \alpha \leq 1$, there is a real $A \leq_{\text{wtt}} \emptyset'$ such that

$$\underline{K}(A) = \alpha \quad \text{and} \quad (\forall Z \leq_{\text{wtt}} A) \underline{K}(Z) \leq \alpha.$$

A Wtt Lower Cone of Non-Integral Dimension

The strategy

- Requirements:

$$R_{\langle e, b \rangle} : Z = \Psi_e(A) \Rightarrow \exists (k \geq j) K(Z \upharpoonright_k) \leq^+ (\alpha + 2^{-b})k$$

where (Ψ_e) is a uniform listing of wtt reduction procedures.

- We can assume each Ψ_e also has a certain (non-trivial) lower bound on the use g_e , because otherwise the reduction would decrease complexity anyway.

A Wtt Lower Cone of Non-Integral Dimension

The strategy

- We construct A inside the Π_1^0 class

$$\mathcal{P} = \{Z : (\forall n \geq n_0) K(Z \upharpoonright_n) \geq \lfloor \alpha n \rfloor\}$$

(This ensures A has dimension at least α .)

- \mathcal{P} is given as an effective approximation through clopen sets P_s .
- We approximate longer and longer initial segments σ_j of A , where σ_j is a string of length m_j , both σ_j, m_j controlled by R_j .

A Wtt Lower Cone of Non-Integral Dimension

The strategy

- Define a length k_j where we intend to compress Z , and let $m_j = g_e(k_j)$.
- Define σ_j of length m_j in a way that, if $x = \Psi_e^{\sigma_j}$ is defined then we compress it down to $(\alpha + 2^{-b_j})k_j$, by enumerating an appropriate request into a Kraft-Chaitin set L .
- The opponent's answer could be to remove σ_j from \mathcal{P} . (σ_j is not of high dimension.)
- In this case, the capital he spent for this removal exceeds what we spent for our request, so we can account our capital against his.
- Of course, usually σ_j is much longer than x . So we will only compress x when the measure of oracle strings computing it is large.

A Wtt Lower Cone of Non-Integral Dimension

An important Lemma

- We assume that \mathcal{P} is effectively approximated by clopen sets P_s .

Lemma

Let \mathcal{C} be a clopen class such that $\mathcal{C} \subseteq P_s$ and $\mathcal{C} \cap P_t = \emptyset$ for stages $s < t$. Then

$$\Omega_t - \Omega_s \geq (\lambda \mathcal{C})^\alpha.$$

A Wtt Lower Cone of Non-Integral Dimension

Combining the strategies R_j

- In the course of the construction, some R_j might have to pick a new σ_j .
- In this case we have to initialize all R_n of lower priority ($n > j$).
- We have to make sure that this does not make us enumerate too much measure into L .
- We therefore have to assign a new length k_n to the strategies R_n .
- For this, it is important to know the use of the reduction related to R_j .

The Turing Case

- It remains an open problem whether there exists a Turing lower cone of non-integral effective dimension.
- This case appears to be much harder. It is, for instance, not even known whether there exists a set of non-integral dimension which does not compute a Martin-Löf random set.

Theorem

There exists computable, non-decreasing, unbounded function f and a set A such that

$$K(A \upharpoonright_n) \geq f(n)$$

and A does not compute a Martin-Löf random set.

[Kjos-Hanssen, Merkle, and Stephan 2004; Reimann and Slaman 2004,]